arXiv:hep-ph/0011119v1 8 Nov 2000

ER/40685/954
UR-1616
November 2000

ELECTROWEAK RADIATIVE CORRECTIONS TO OFF-SHELL W-PAIR PRODUCTION [*]

ANSGAR DENNER
Paul Scherrer Institut
CH-5232 Villigen PSI, Switzerland

STEFAN DITTMAIER
Theoretische Physik, Universität Bielefeld
D-33615 Bielefeld, Germany

MARKUS ROTH
Institut für Theoretische Physik, Universität Leipzig
D-04109 Leipzig, Germany

DOREEN WACKEROTH
Department of Physics and Astronomy, University of Rochester
Rochester, New York 14627, USA

We briefly describe the RacoonWW approach to calculate radiative corrections to $e^+e^- \to WW \to 4f$ and present numerical results for the total W-pair production cross section at LEP2.

1. Introduction

At LEP2, for the first time, a precise direct measurement of the triple gauge-boson couplings $(\gamma, Z)WW$ is performed[1], allowing to test the non-abelian structure of the Standard Model of electroweak interactions (SM). Moreover, a measurement of the W-boson mass at LEP2 with a precision of 35 MeV is within reach[1], which will not only provide a further precisely known SM input parameter but will also help to considerably improve the present indirect bounds on the Higgs-boson mass[2]. To match the experimental precision at LEP2 and a future Linear e^+e^- Collider, the theoretical predictions for the cross sections to $e^+e^- \to WW \to 4f$ need to be of $\mathcal{O}(1\%)$ accuracy and better[3]. This requires to go beyond the inclusion of universal corrections such as the running of the electromagnetic coupling, corrections to the ρ parameter, the Coulomb singularity, and leading photonic corrections, as described in Ref.4. The theoretical uncertainty of predictions which only include universal radiative corrections is estimated[4] to be of $\mathcal{O}(1$–$2\%)$ for the total W-pair production cross section, σ_{WW}, at LEP2 energies and is expected to be even larger for differential cross sections. So far, no complete calculation of the electroweak $\mathcal{O}(\alpha)$ corrections to the processes $e^+e^- \to 4f$ exists. Such a calculation is not only enormously complex but also poses severe theoretical problems with gauge invariance connected to the instability of the gauge bosons. Nevertheless, there are efforts in this direction[5]. At present, all predictions for $e^+e^- \to WW \to 4f$, which go beyond universal radiative corrections, rely on a Double-Pole-Approximation (DPA): radiative corrections are taken into account only to those contributions that are enhanced by two resonant W propagators. Different versions of the DPA are implemented in the Monte Carlo generators RacoonWW[6] and YFSWW3[7], and the results of a tuned comparison at LEP2 and LC energies are presented in Refs.3 and 8, respectively. As a result of this comparison, supported by a tuned comparison with the semi-analytic calculation of Ref.9 and a detailed study of the intrinsic uncertainty of the DPA with RACOONWW, the four-fermion working group of the LEP2MC workshop[3] assigned a theoretical uncertainty to the predictions for σ_{WW} by RACOONWW and YFSWW3 of 0.4% at energies between 200 and 500 GeV. In the following we give a brief description of the RACOONWW approach.

---

[*] Talk given at the DPF 2000 meeting, Columbus, OH, August 9-12, 2000.

2. Electroweak Radiative Corrections to $e^+e^- \to WW \to 4f$

The complete $\mathcal{O}(\alpha)$ corrections to off-shell W-pair production in DPA as implemented in RACOONWW can be written as follows[6]:

$$\mathrm{d}\sigma_{\mathrm{WW}} = \mathrm{d}\sigma^{e^+e^-\to 4f}_{\mathrm{Born}} + \mathrm{d}\sigma^{e^+e^-\to WW\to 4f}_{\mathrm{virt,finite,DPA}} + \mathrm{d}\sigma^{e^+e^-\to 4f}_{\mathrm{virt+real,sing}} + \mathrm{d}\sigma^{e^+e^-\to 4f\gamma}_{\mathrm{finite}}. \qquad (1)$$

Here $\mathrm{d}\sigma^{e^+e^-\to 4f}_{\mathrm{Born}}$ is the full lowest-order cross section to $e^+e^- \to 4f$ and $\mathrm{d}\sigma^{e^+e^-\to 4f\gamma}$, which describes real photon radiation away from IR and collinear singularities, is the full lowest-order cross section to $e^+e^- \to 4f\gamma$ as described in Ref.10. The cross section for real photon radiation is rendered finite in two different ways: by using phase-space slicing or alternatively a subtraction method[11]. The exact $4f$ and $4f+\gamma$ phase spaces for massless fermions are used throughout. The IR-finite sum of virtual, soft and collinear photonic one-loop corrections is denoted by $\mathrm{d}\sigma^{e^+e^-\to 4f}_{\mathrm{virt+real,sing}}$. Apart from finite terms, it contains only collinear singularities associated with the initial state, i.e. leading logarithms of the form $\ln(s/m_e^2)$, at least for inclusive enough observables. Since the contribution $\mathrm{d}\sigma^{e^+e^-\to 4f}_{\mathrm{virt+real,sing}}$ is not treated in DPA, those logarithms are treated in our approach exactly. In RACOONWW, the DPA is only applied to the finite part of the virtual corrections, denoted by $\mathrm{d}\sigma^{e^+e^-\to WW\to 4f}_{\mathrm{virt,finite,DPA}}$ including the full set of factorizable and non-factorizable virtual $\mathcal{O}(\alpha)$ corrections. The factorizable contribution comprises those virtual one-loop corrections that can either be assigned to the W-pair production or to the W-decay subprocesses and, thus can be deduced from the known results for on-shell W-pair production and W decay. The non-factorizable corrections connect the production and decay processes. In DPA, they comprise all doubly-resonant contributions that are not already included in the factorizable corrections. Beyond $\mathcal{O}(\alpha)$, RACOONWW includes soft-photon exponentiation and leading higher-order initial-state radiation up to $\mathcal{O}(\alpha^3)$ via the structure-function approach[4]. The leading higher-order effects from $\Delta\rho$ and $\Delta\alpha$ are taken into account by using the G_μ scheme. QCD corrections are taken into account either by a multiplicative factor $(1 + \alpha_s/\pi)$ for each hadronically decaying W boson or as complete QCD corrections to the CC03 diagrams.



Fig. 1. The total W-pair production cross section and the ratio of the data over the RacoonWW prediction at LEP2, as given by the LEPEWWG[2].

Numerical results from RACOONWW at LEP2 energies for some interesting observables, such as invariant-mass distributions of the W decay products and angular distributions, are provided in Refs.3,6,12. In Fig.1, we show the comparison of the RACOONWW prediction for the W-pair-production cross section with LEP2 data. From a comparison of RACOONWW with GENTLEv2.00[13], for instance, the neglect of non-universal $\mathcal{O}(\alpha)$ corrections is known to induce a 2.5% shift[3] to higher values of σ_{WW} which is disfavoured by the data. The good agreement of LEP2 data with the state-of-the-art Monte Carlo generators RACOONWW and YFSWW3 reveals evidence of non-leading electroweak corrections beyond the level of universal effects.

1. A. Gurtu, talk given at ICHEP 2000, Osaka, July 17 - August 2, 2000.
2. D. Abbaneo *et al.*, CERN-EP-2000-016 and lepewwg.web.cern.ch/LEPEWWG/.
3. M. Grünewald *et al.*, in *LEP2MC workshop* (CERN, 1999/2000), hep-ph/0005309.
4. W. Beenakker *et al.*, in *Physics at LEP2*, eds. G. Altarelli *et al.* (Report CERN 96-01, Geneva, 1996, Vol. 1, p. 79), hep-ph/9602351.
5. A. Vicini, *Acta Phys. Pol.* **B29**, 2847 (1998).
6. A. Denner *et al.*, *Nucl. Phys.* **B587**, 67 (2000).
7. S. Jadach *et al.*, *Phys. Lett.* **B417**, 326 (1998); *Phys. Rev.* **D61**, 113010 (2000); hep-ph/0007012.
8. A. Denner *et al.*, in *MRST 2000*, ed. C.R. Hagen (Melville, NY, 2000), hep-ph/0007245.
9. W. Beenakker *et al.*, *Nucl. Phys.* **B548**, 3 (1999).
10. A. Denner *et al.*, *Nucl. Phys.* **B560**, 33 (1999).
11. S. Dittmaier, *Nucl. Phys.* **B565**, 69 (2000); M. Roth, PhD thesis, hep-ph/0008033.
12. A. Denner *et al.*, *Phys. Lett.* **B475**, 127 (2000).
13. D. Bardin *et al.*, *Phys. Lett.* **B308**, 403 (1093) and *Comput. Phys. Commun.* **104**, 161 (1997).